



Ministry of Health
& Family Welfare
Government of India



Introduction of **Rotavirus Vaccine** in the Universal **Immunization** Programme

Frequently Asked Questions

For Health Workers



Given Orally 5 drops

Immunization Division

March 2019



Be Wise!
**Get your child
fully immunized**

Frequently Asked Questions on Rotavirus Vaccine Introduction For Health Workers



1. What is Rotavirus?



Rotavirus is a highly contagious virus. It is the most common organism that causes diarrhea in children which may lead to hospitalization and death.



2. What are the clinical features of Rotavirus diarrhea?



The clinical features of Rotavirus diarrhea may vary from transient loose stools to severe diarrhea with vomiting. If not treated adequately, it may result in dehydration, electrolyte imbalance, shock and death.

Rotavirus diarrhea has an incubation period of 1 to 3 days. It presents usually with sudden onset of watery stools, often accompanied by fever and vomiting. Sometimes there may be abdominal pain. The diarrhea and associated symptoms may last for 3 to 7 days.



3. Is Rotavirus diarrhea serious?



Rotavirus accounts for 40% of hospitalization due to diarrhea amongst children in India. It is responsible for nearly 32.7 lakh outpatient visits, 8.72 lakh hospitalizations and 78000 deaths annually in India of which 59,000 occur in the first two years of life.¹

¹John J, Sarkar R, Muliyl J, Bhandari N, Bhan MK, Kang G. Rotavirus gastroenteritis in India, 2011-2013: Revised estimates for disease burden and potential impact of vaccines. *Vaccine* 2014;32 Suppl.1: A5-9



4. Which children are at risk of getting the Rotavirus diarrhea?



The risk of Rotavirus infection continues to persist in all settings. The proportion of diarrhea caused by Rotavirus does not vary widely between developed and developing countries.

Approximately 50% of Rotavirus-associated deaths occur in the first year of life and about 75% occur in the first two years of life.

There is a high risk of dehydration and death due to diarrhea in undernourished children, if not treated adequately.



5. What are the modes of transmission of the Rotavirus?



Rotavirus is a highly contagious virus. Rotavirus transmission occurs primarily by the faeco-oral route directly from person to person, or indirectly via contaminated fomites and the virus can live for hours on hands and even longer on hard surfaces.



6. In which season does Rotavirus diarrhea occur most?



Rotavirus infection and diarrhea can occur throughout the year, most commonly seen in winter season.



7. How is Rotavirus diarrhea diagnosed?



Rotavirus diarrhea cannot be clinically distinguished from diarrhea due to other causes. Rotavirus diarrhea can be diagnosed in stool specimen, using laboratory tests.



8. What is the treatment of Rotavirus diarrhea?



There is no specific treatment for Rotavirus diarrhea. The treatment of Rotavirus diarrhea is similar to other acute diarrheas, using ORS for correcting fluid and electrolyte imbalance and zinc for 14 days. Severe dehydration may require hospitalization for treatment with intravenous (IV) fluids.



9. Once a child has had Rotavirus diarrhea, can she/he get it again?



Yes, the child can get repeated infection with Rotavirus. The subsequent infections may be milder in nature.



10. How can Rotavirus diarrhea be prevented?



Rotavirus vaccine is the only specific prevention against Rotavirus diarrhea. The vaccine is effective in reducing the hospitalizations and deaths due to diarrhea in children.

General measures to prevent diarrhea like good hygiene, frequent hand washing, safe water and safe food consumption, exclusive breastfeeding and vitamin A supplementation reduces the risk of Rotavirus infection, but are not enough to control the spread of the disease.



11. Will vaccination with Rotavirus vaccine prevent all diarrheas?



No. Please remember that diarrhea is caused by many organisms. Rotavirus vaccine is effective in preventing diarrhea due to Rotavirus only, which is one of the leading cause of hospitalizations and deaths due to diarrhea in children. The child may still get diarrhea due to germs and other reasons, even after receiving the vaccine.



12. What will be the presentation of the Rotavirus vaccine?



Rotavirus vaccine is an oral, liquid vaccine and supplied in a five dose vial along with one dropper per vial.



13. What is the appearance of Rotavirus vaccine?



The Rotavirus vaccine is generally pink in colour and may sometimes change to orange or light yellow. This change in colour does not impact the quality of the vaccine.




14. How and when is the Rotavirus vaccine given?



The dose of this liquid Rotavirus vaccine currently being supplied under UIP is 5 drops (0.5 ml) given orally. Rotavirus vaccine will be administered to all infants at 6, 10 and 14 weeks along with other vaccines in routine immunization schedule as per the preferred sequence given below:

- **At 6 weeks:** OPV (2 drops oral) - Rotavirus vaccine (5 drops oral) – fIPV (0.1 ml intradermal) – PCV (0.5 ml IM)*-Pentavalent vaccine (0.5 ml IM)
- **At 10 weeks:** OPV (2 drops oral) - Rotavirus vaccine (5 drops oral) - Pentavalent vaccine (0.5 ml IM)
- **At 14 weeks:** OPV (2 drops oral) - Rotavirus vaccine (5 drops oral) – fIPV (0.1 ml intradermal) – PCV (0.5 ml IM)*-Pentavalent vaccine (0.5 ml IM).

** Wherever applicable*



15. Is a booster dose required for Rotavirus vaccine?



No booster dose of Rotavirus vaccine is recommended. Only three doses at 6, 10 and 14 weeks are required to complete the vaccination schedule for a child.



16. Does breastfeeding need to be withheld for some time before or after vaccination?



The infant can be breastfed before or after vaccination.



17. What are the clinical contraindications for Rotavirus vaccination?



The clinical conditions where Rotavirus vaccine must not be administered to the infant are:

- a. Known or documented allergic reaction to the vaccine
- b. History of documented intussusception or abdominal surgery or intestinal malformation
- c. Known case of immunodeficiency.

Q

18. If the infant has minor illness, can Rotavirus vaccine be given?

A

If the infant has minor illnesses like mild fever, upper respiratory infection etc., Rotavirus vaccine can be given. If the infant is well enough to have the other routine vaccinations, then Rotavirus vaccine can also be given. There's no reason to postpone the vaccination unless the infant is seriously ill, having high grade fever, severe diarrhea or vomiting.

Q

19. Where should the Rotavirus vaccine be recorded on MCP card?

A

Rotavirus vaccine should be recorded in the space provided beside Rota-1, Rota-2 and Rota-3 on MCP cards.

BIRTH	1 1/2 MONTHS	2 1/2 MONTHS	3 1/2 MONTHS	3 MONTHS
Expected date of delivery / /	Next Vaccination Date: / /	Next Vaccination Date: / /	Next Vaccination Date: / /	Next Vaccination Date: / /
DATE GIVEN (mm/dd/yyyy)	DATE GIVEN (mm/dd/yyyy)	DATE GIVEN (mm/dd/yyyy)	DATE GIVEN (mm/dd/yyyy)	DATE GIVEN (mm/dd/yyyy)
OPV-0 / /	OPV-1 / /	OPV-2 / /	OPV-3 / /	MR-1 / /
Hep B give within 24h of birth / /	Penta-1 / /	Penta-2 / /	Penta-3 / /	JE-1 / /
BCG / /	Rota-1 / /	Rota-2 / /	Rota-3 / /	Vitamin A-1 / /
/ /	PCV-1 / /	/ /	PCV-2 / /	PCV booster / /
/ /	IPV-1 / /	/ /	IPV-2 / /	/ /
/ /	/ /	/ /	/ /	/ /

Neonatal Care

Please remember:

- Keep the child warm.
- Start breastfeeding within 1 hr after birth
- Feed the baby only mother's milk.
- Do not bathe the child for the first 48 hours
- Keep the cord dry
- Keep the child away from sick people
- Special care if child > 2.5 kg at birth.

Danger signs:

Contact your health worker if the baby:

- is sucking weakly or refuses to breast feed
- is unable to cry or has difficulty in breathing
- Has yellow palms and soles
- Has fever or a cold to touch
- Has blood in stools or convulsions.
- Is lethargic or unconscious

Congratulations! Your child is vaccinated for the 1st year of life.

Q

20. Is it safe to give multiple vaccinations to a child in one visit?

A

Yes. It is safe to give multiple vaccinations as per national immunization schedule to a child in one visit.

Q

21. What is the maximum age limit for giving the first dose of Rotavirus vaccine?



A

The upper age limit for giving the first dose of Rotavirus vaccine is one year of age. If the child has received first dose of Rotavirus vaccine by one year of age, subsequent 2nd and 3rd dose of the vaccine should be given at an interval of 4 weeks between two doses to complete the course.

Q

22. What are the similarities and differences between oral liquid Rotavirus vaccine and oral Polio vaccine?

A

S.No	Feature	Oral Rotavirus Vaccine	Oral Polio Vaccine
Similarity			
1.	Colour of vaccine	Generally pink in colour, may change to orange or light yellow	Light pink or yellowish orange.
2.	Mode of administration	Oral	Oral
Differences			
3.	Colour of dropper	Pink	Transparent
4.	Taste	Sweet	Bland to Bitter
5.	Dose	5 drops	2 drops
6.	Schedule	6,10 & 14 week of age	OPV zero dose for institutional deliveries. Primary doses at 6,10 & 14 weeks; Booster dose at 16-24 months of age.
7.	VVM	On Cap 	On label 

Q

23. Should Rotavirus vaccine be started in children who have already received first dose of OPV and Pentavalent vaccine?

A

No. During the initial period of Rotavirus vaccine introduction, only the infants coming for the first dose of OPV and Pentavalent vaccine will be started with Rotavirus vaccine. These children will be given 2nd and 3rd doses in subsequent visits as per the schedule.



24. What if a child misses the first dose of Rotavirus vaccine or comes late for first dose vaccination?



If the child comes after 6 weeks of age for first dose of recommended UIP vaccines, the first dose of Rotavirus vaccine can be given along with first doses of OPV and Pentavalent vaccine upto one year of age. The upper age limit for starting Rotavirus vaccination is one year of age.

If the child has received first dose of Rotavirus vaccine by one year of age, subsequent 2nd and 3rd dose of the vaccine to be given with an interval of 4 weeks between two doses to complete the course.



25. Will Rotavirus vaccine be given in Intensified /Mission Indradhanush?



Yes, Rotavirus vaccine will be given in Intensified /Mission Indradhanush following the same guidelines as for Routine Immunization.



26. Should Rotavirus vaccine be given to a child coming from a state which has not yet introduced Rotavirus vaccine in its schedule?



Yes, Rotavirus vaccine should be given to a child irrespective of the state where she/he comes from, as per the current guidelines.



27. Should the liquid Rotavirus vaccine be given to a child who has received one or two doses of an oral freeze-dried Rotavirus vaccine available under UIP in another state?



If a child starts the schedule with the oral freeze dried Rotavirus vaccine then the child should preferably complete the schedule using the same oral freeze dried vaccine. However, in case of inter-state migration, vaccination should not be deferred or denied because the Rotavirus vaccine used for the previous dose (s) is unknown or is different from the Rotavirus vaccine available in the state where the child's family has migrated. The RVV schedule will be completed using the Rotavirus vaccine available in that state under the UIP.



28. What should be done if a child has received one or two doses of Rotavirus vaccine in private facility?



Please remember that there are four Rotavirus vaccines products available in the market. If a child starts the schedule with the Rotavirus vaccine product "A" then the child should preferably complete the schedule using the same Rotavirus vaccine product 'A'. However, vaccination should not be deferred or denied because the Rotavirus vaccine product used for the previous dose (s) is unknown or is different from the Rotavirus vaccine product available under UIP in that state. The subsequent doses will be administered using the Rotavirus vaccine available under UIP in that state to complete the RVV schedule.



29. What if a child who has received first dose of RVV and has missed second or third dose of Rotavirus vaccine?



The child can be given due doses of Rotavirus vaccine (2nd and 3rd dose) at a minimum interval of 4 weeks between two doses.



30. What is to be done if the child is given Rotavirus vaccine first before OPV?



No problem. If the child is given Rotavirus vaccine first, give OPV after that.



31. Is the Rotavirus vaccine schedule for prematurely born babies different?



No. The doses of Rotavirus vaccine are to be administered as per the age calculated from date of birth, irrespective of the age of gestation when the child was born.



32. Which government health facilities will provide Rotavirus vaccine?



Rotavirus vaccine will be provided free of cost through routine immunization sessions under UIP. The vaccine will be provided at all government health facilities including hospitals, medical colleges, urban dispensaries, PHCs, CHCs, sub centers and outreach session sites. It cannot be given in house-to-house visits.



33. What should be done if Rotavirus vaccine is not available at immunization session site?



The child should be given all available vaccines for which she/he is eligible. Rotavirus vaccine should be given to the child in the next immunization session.



34. Is there any side effect of the Rotavirus vaccine?



Rotavirus vaccine is a safe vaccine. However, mild and transient symptoms including vomiting, diarrhea, cough, runny nose, fever, irritability and rash may be associated with Rotavirus vaccination.

A rare adverse event known as intussusception (one part of bowel folding/telescoping into another) has been reported after Rotavirus vaccine. As per the WHO position paper (2013), "in some but not all settings, post-marketing surveillance of both currently available Rotavirus vaccines has detected a small increased risk of intussusception (about 1–2/100 000 infants vaccinated) shortly after the first dose."

Such children present with severe pain abdomen (excessive crying) and repeated episodes of vomiting, blood in stool and dehydration. Such children need to be referred immediately to hospital for appropriate management. Also, in such cases, inform DIO and report this AEFI case as per guidelines

It is important to know that Intussusception also occurs in children without Rotavirus vaccine.

The Rotavirus vaccine offers tremendous benefits by protecting infants and children from Rotavirus disease. The risk of adverse events after Rotavirus vaccination is much lower than the risk of severe Rotavirus disease in unvaccinated children. Hence, Rotavirus vaccine is strongly recommended to prevent Rotavirus disease in infants and young children.



35. Will the Rotavirus vaccine vials have vaccine vial monitor (VVM)?



Yes. Rotavirus vaccine vials will have a vaccine vial monitor (VVM) in the cap of the vaccine vial. If the VVM reaches the discard point, do not use the vaccine.

As long as the color of inner square is lighter than the colour of the outer ring, the vaccine can be used. As soon as the colour of the central square is the same colour as the ring or of a darker colour than the ring, the vial should be discarded.

How to read a VVM

- ✓ Vaccine OK
- ✓ Vaccine OK but use first
- ✗ Do not use the vaccine
- ✗ Do not use the vaccine



36. At what temperature should Rotavirus vaccine be stored?



Rotavirus vaccine should be stored between -15°C and -25°C in GMSDs, State, Regional and District stores in the walk-in freezers (WIFs) or deep freezers (DFs). Cold chain points below district level should store RVV at +2°C to +8°C, in ice-lined refrigerators (ILRs). In the ILR, the RVV should be stored at or above BCG level.

The droppers for administration of the vaccine are to be stored at room temperature as freezing can cause the droppers to crack. They are to be supplied to the immunization session sites` along with other dry supplies outside the vaccine carrier .

✓ To be used

Opened vials of Hep. B, DPT, TT/Td, Pentavalent, OPV, IPV and PCV provided;

- a. VVM is intact and in usable stage
- b. Date and Time of opening is mentioned on the vial
- c. Vial is within 28 days of opening
- d. Date of expiry has not reached/crossed
- e. The vaccine vial septum is not contaminated

✗ Not to be used

1. Opened vials of BCG, MR, JE and RVV
2. All empty vaccine vials
3. VVM is not in usable stage
4. Date and time of opening is not mentioned
5. 28 days of opening of vial have passed

*These vials should be discarded after 48 hours or before the next session, whichever is earlier as per the CPCB guidelines.



37. How should Rotavirus vaccine be transported between different cold chain points?



Like other UIP vaccines, Rotavirus vaccine should be transported in cold boxes with conditioned ice-packs..



38. How should Rotavirus vaccine be transported to session sites?



Rotavirus vaccine should be transported to session sites along with other vaccines in a vaccine carrier with four conditioned ice packs. The droppers are not to be kept inside the vaccine carrier, and should be sent separately along with other dry supplies.



39. Does Open Vial Policy apply to Rotavirus vaccine?



The open vial policy is **NOT** applicable to Rotavirus vaccine, similar to JE, MR and BCG vaccines.

A Rotavirus vaccine vial can be used up to a maximum of **4 hours** after opening. It is mandatory to write the date and time of opening on the vaccine vial. All partially used vaccine vials should be sent back to cold chain point for disposal as per biomedical waste management guidelines.



40. What is the maximum time till when an opened Rotavirus vaccine vial can be used?



The Rotavirus vaccine vial once opened can be used up to maximum of **4 hours**.



41. What should you do if you find a frozen Rotavirus vaccine vial?



You can use the vial after it liquifies.

Rotavirus vaccine is stored between -15°C and -25°C at district level and above and between +2°C and +8°C at the sub district level. The vaccine needs to be liquified before it is used.



42. What If the child spits out the Rotavirus vaccine or vomits immediately after having it?



Repeat the dose. In case an incomplete dose is administered (the infant spits out or regurgitates most of the vaccine), repeat the dose in the same vaccination visit. To prevent spitting, please put the drops towards the inner cheek. Avoid putting the drops over the tongue.



43. Where should the Rotavirus vaccine vial be kept at the immunization site?



Please place the opened Rotavirus vaccine vial on the ice pack kept outside during the session. As per RI guidelines, the health worker is expected to take out one ice pack at the session site and use the same for placing vaccines that are not freeze sensitive. On ice pack- BCG and MR (place them in the wells on ice pack), OPV, JE and Rotavirus vaccine should be placed on the surface of the ice pack. Remember – vials of freeze sensitive vaccines including IPV, Td, DPT, Pentavalent and PCV vaccines should never be kept on the ice pack.



Steps in administration of Rotavirus vaccine



Vaccine Vial & dropper



Pull out the aluminum seal along the indicated mark



Tear off as shown to remove aluminum seal



Tear off as shown to remove aluminum seal



Vaccine vial without aluminum seal



Pull out the rubber stopper



7 Connect the dropper firmly to the vial. Write date and time of opening the vial.



8 Open the dropper cap



9 Position dropper at 45° angle. Administer 5 drops into the mouth of the baby 0.5mL = 5 drops*
** The dropper should not touch the mouth of the baby.*



10 Once opened the multi dose vial should be kept on conditioned icepack (2° - 8°) and cannot be used beyond 4 hours.



World Health
Organization

unicef



IMMUNIZATION
TECHNICAL SUPPORT UNIT
Ministry of Health & Family Welfare

